

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

Reserve
A423.9
F764

U. S. DEPT. OF AGRICULTURE

U.S. DEPARTMENT OF AGRICULTURE FOREST PEST LEAFLET 75

Forest Service

FEB 8 - 1963

January 1963

~~Beech Bark Disease~~

By Alex L. Shigo¹

American beech (*Fagus grandifolia* Ehrh.) is an important component of the northern hardwood forests, which occupy about 29 percent of the commercial forest land in the New England and Middle Atlantic States. Even though beech wood has desirable qualities closely akin to those of sugar maple and yellow birch, for a long time beech was discriminated against for lumber and other high-quality products because it was difficult to season and process. But now, as improved techniques have made it easier to season and process beech, this species has become quite acceptable in the lumber markets; and the wood is used increasingly for a variety of products, such as furniture, flooring, handles, dowels, shuttles, and spools. In recent years, more beech has also been used as pulpwood, and with new processing techniques it may become even more valuable for this purpose.

The growing of large, high-quality beech is complicated by the beech bark disease. This disease has killed more than half the beech in some stands, and many of the surviving trees hold little promise for future production of quality material.

Very little research has been done on the beech bark disease since 1934, possibly because of the old prejudices against beech wood and its low standing in the lumber markets. Most reports on the disease in the past 25 years have done little more than tell how rapidly the disease was spreading.

History and Distribution

Accounts from Europe indicate that beech bark disease was killing beech trees before 1849. The obvious scale insect (*Cryptococcus fagi* (Baer.)) on beech was considered the cause until 1914, when it was learned that the fungus *Nectria ditissima* infected the trees infested by the insects.

The insect and probably the associated fungus were introduced into North America on ornamental beech trees. The first reported outbreak of the disease was in 1920 in Nova Scotia, Canada, about 30 years after the insect was known to be present in the area of Halifax. In 1929 the insect was found on ornamental beech trees in the Arnold Arboretum in Boston. At the same time a *Nectria* was noted on the trees dying after scale infestations in Nova Scotia. By 1932 the disease was known throughout the beech areas of the Maritime Provinces in Canada, in localized areas of eastern and south-central Maine, and in eastern Massachusetts.

Most of the research work on the disease in North America was done at that time. Since then the disease has continued to spread southward and westward to beech areas throughout the New England States and eastern New York. Recently the insect has been reported from areas in central New York and northern Pennsylvania. Continued spread of the disease appears likely.

¹ Forest pathologist, Northeastern Forest Experiment Station, Forest Service, U.S. Department of Agriculture. Mr. Shigo is stationed at Laconia, N.H.

The Cause

American beech (*Fagus grandifolia*), European beech (*F. silvatica*), and all varieties of these species are hosts in varying degrees to the bark-killing fungus, *Nectria coccinea* var. *faginata* Lohman, Watson, and Ayres, which infects through minute wounds and cracks in the bark made by the feeding tubes of the beech scale insect, *Cryptococcus fagi*. Although a fungus and an insect are associated in this disease, the insect was observed first and has been studied in greater detail than the fungus.

In Europe the *Nectria* first observed associated with *Cryptococcus fagi* was called *Nectria ditissima*, and is now referred to as *N.*

galligena. In North America the fungus associated with *C. fagi* is thought to be a variety of a common weakly parasitic species, *N. coccinea* var. *faginata*. Several other species of *Nectria* have been isolated from some trees showing symptoms of this disease.


The Insect

Cryptococcus fagi is a soft-bodied scale insect. At maturity it is yellow, elliptical, $\frac{1}{2}$ to 1 mm. long, has reddish-brown eyes, a stylet approximately 2 mm. long, rudimentary antennae and legs, and numerous minute glands that secrete a white, woollike wax over the entire body (fig. 1).



F-502240

FIGURE 1.—A section of beech bark covered with the white wax secretion of the beech scale.



F-502241

FIGURE 2.—Second stage larvae of *Cryptococcus fagi*. The wax was removed before photographing. The larvae are 0.4 to 0.6 mm. long.

Reproduction is parthenogenetic, males being unknown. Beginning in midsummer, the insect deposits pale yellow eggs on the bark in strings of four to eight, attached end to end. Depending on temperature, the eggs usually begin to hatch in late summer, and hatching continues until early winter.

The larvae emerge from the eggs with well-developed legs and antennae. The larvae either remain stationary under the females that died after egg deposition, migrate to cracks and other areas where they will be protected, fall to the ground where death is imminent, or establish themselves on other trees after being disseminated by various agents. After the insect becomes stationary, it forces its tubular stylet into the bark. It is then a second-instar nymph, without legs and covered with woollike wax (fig. 2). The insect hibernates in this stage on the bark and moults in the spring to become an adult female.

The Fungus

Nectria coccinea var. *faginata* is a microfungus that produces several types of spores. The conspicuous bright red, lemon-shaped, fruiting bodies (perithecia) (fig. 3) seen on the trees in clusters of as many as 40 are filled with elongate sacs, each containing 8 spores. These spores result from a sexual process, and their production constitutes the perfect stage of the fungus.

The perithecia mature in the fall, and the spores are forced out only when they have been sufficiently moistened. After these spores dry they appear as white dots on the tips of the perithecia. Perithecia on the dead bark continue to produce viable spores the following year.

The other types of spores do not follow a sexual process, and this is called the imperfect stage. Small white cushions of asexual spores



F-502242

FIGURE 3.—Fruiting bodies (perithecia) of *Nectria coccinea* var. *faginata*. They may be in clusters up to 40 or in a single line. The perithecia are approximately 0.3 x 0.25 mm.



F-502243

FIGURE 4.—The imperfect stage of *Nectria coccinea* var. *faginata*, growing from fungus tissue in the bark. The tufts of spore-bearing branches, 1 to 4 mm. broad, can easily be mistaken for isolated colonies of the scale insect.

(fig. 4) frequently burst through the bark before the perithecia appear. These can easily be mistaken for small isolated colonies of the scale insect. The asexual spores range from single-celled oval spores to eight-celled sickle-shaped spores. These spores are produced in a dry head, well suited for wind dissemination. The imperfect stage can be found on the trees from midsummer until fall. Because of the variability of the fungi in the *Nectria coccinea* group, there is still confusion over the correct name of the *Nectria* that follows the scale infestation.

Symptoms and Course of the Disease

First symptoms are isolated dots of white "wool" that appear on roughened areas of the bark, under branches, and in the lenticels. Eventually the entire bole of the tree may be covered with the waxy material secreted by the scale insects as they increase in numbers (fig. 1). Heavy infestations feeding on the liquids in the bark cells can greatly weaken a tree.

On some trees a red-brown liquid exudes from dead spots near areas heavily infested with the insects.

This seeps down the tree, forming a slime-flux or a brown spot. There is no satisfactory explanation for this: the most common theory is that the slime-flux is a secondary effect of the necrosis produced by the feeding of the insects on the cells and that other fungi infect these areas, giving them a foul odor. Frequently perithecia of the *Nectria* are later found around these dead spots. The dead areas may extend into the sapwood. These spots are usually delimited by callus tissue.

Areas devoid of the scale insects or patches of black wool indicate one of the first places killed by *Nectria*. The insects cannot live on dead tissues; and, as they die, a black fungus frequently grows over them. The necessary time and conditions from the first insect infestation to the first fungus infection are not understood, but some are of the opinion that 3 to 6 years are necessary under optimum conditions for infection to be made obvious by the production of the bright red perithecia.

The fungus may infect large areas on some trees, completely girdling them. The leaves that do come out in the spring do not mature, giving the crowns an open appearance. The leaves turn yellow and usually remain on the tree during the summer season. The chlorotic crowns are typical of trees dying from a water deficiency.

Frequently the fungus infects only narrow strips on the bole, and the subsequent symptoms differ from those of trees that have been girdled. Callus tissue forms around these cankers and the bark becomes very roughened (fig. 5). Parts of the crown become chlorotic and die. Small cankers may be walled off from the sapwood by callus tissue (fig. 6).

Other insects and wood-rotting fungi quickly invade the large necrotic areas. Many of the trees that




F-502244

FIGURE 5.—*Fomes igniarius* and *Hypoxylon* sp. growing on an area killed by the *Nectria*. Callus tissue has caused the bark to crack along the large lesion. Trees that are not girdled by the *Nectria* often live a long time.

are partially girdled remain alive, in a weakened state, for many years, while others are broken by the wind. Species of *Hypoxylon* that decay sapwood are the first to invade the necrotic areas. Several species of ambrosia beetles facilitate the ingress of other fungi.

Vigorous trees, free of the disease, are often found in heavily infected areas. As some investigators have pointed out, this suggests that some trees are more resistant to the disease than others (fig. 7).



F-502245

FIGURE 6.—The craterlike scars indicate where small isolated *Nectria* cankers were walled off by callus tissue. The disease has caused only moderate injury to these trees.



F-502246

FIGURE 7.—The beech on the left is free of scale and *Nectria*; the one on the right has been recently killed by the disease. This suggests that certain trees are resistant to the disease.

Control

The beech scale insect has very few natural enemies and leads a fairly unmolested life. Air temperatures of -35° F. and lower are lethal to those insects not protected by snow. A ladybird beetle,

Chilocorus stigma, feeds on the scale; but its effect on the scale population is thought to be negligible.

The fungus *Gonatorrhodiella highlei* is parasitic on the *Nectria*. This mycoparasite is often found covering large areas of the perfect and imperfect forms of the *Nectria*.

The effect of the fungus on *Nectria* is not known.

Several insecticides such as DDT and lime-sulfur are effective against the scale. At present it is highly impractical to consider spraying trees grown for timber, unless they are in areas where intensive cultural measures would be practicable. Insecticides may be used on trees grown for their esthetic value.

Conflicting reports on the effect of stand density on the progress of the disease have been presented. A program of timely salvage cuttings is the only way presently known to keep the losses from this disease to a minimum.

Caution: DDT is poisonous. Store it in plainly labeled containers away from all food. Follow directions and heed precautions given by the manufacturer.

References

- STUDIES OF PYRENOAMYCETES: IV. *NECTRIA* (PART I). C. Booth. Commonwealth Mycol. Inst. Mycol. Paper 73, 115 pp. 1959.
- NOTES ON THE BEECH SCALE, *CRYPTOCOCCUS FAGI* (BAER.) DOUGL., NEW ENGLAND. R. C. Brown. Jour. Econ. Ent. 27: 327-334. 1934.
- TIMELY SALVAGE CAN REDUCE LOSSES FROM BEECH SCALE-*NECTRIA* ATTACK. D. Crosby and J. C. Bjorkbom. U.S. Forest Serv. Northeast. Forest Expt. Sta. Forest Res. Note 82, 4 pp. 1958.
- THE BEECH BARK DISEASE, A *NECTRIA* DISEASE OF *FAGUS*, FOLLOWING *CRYPTOCOCCUS FAGI* (BAER.). J. Ehrlich. Canad. Jour. Res. 10: 593-692. 1934.
- IDENTITY AND HOST RELATIONS OF *NECTRIA* SPECIES ASSOCIATED WITH DISEASES OF HARDWOODS IN EASTERN STATES. M. L. Lohman and A. J. Watson. Lloydia 6: 77-108. 1934.
- ANGREB OF *CRYPTOCOCCUS FAGI*, *NECTRIA GALLIGENA* OG ANDRE PARASITER PAA BØG I DANMARK 1939-43. M. Thomsen, N. F. Buchwald, and P. Hauberg. Forstl. Forsgsv. i Danmark. Beret. 18 (157): 97-326. (English summary.) 1949.